A spectral survey of CH_3CCH in the Hot Molecular Core G331.512-0.103

JULIA C. SANTOS D^{1,2} LEONARDO BRONFMAN,³ EDGAR MENDOZA D^{1,4} JACQUES R. D. LÉPINE D^{1,1} NICOLAS U. DURONEA,^{5,6} MANUEL MERELLO ³ AND RICARDO FINGER³

¹Instituto de Astronomia, Geofísica e Ciências Atmosféricas; Universidade de São Paulo, Rua do Matão, 1226, São Paulo, SP, Brazil ²Laboratory for Astrophysics, Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands

³Departamento de Astronomía; Universidad de Chile, Casilla 36-D, Santiago de Chile, Chile

⁴Observatório do Valongo; Universidade Federal do Rio de Janeiro, Ladeira do Pedro Antônio, 43, Rio de Janeiro, RJ, Brazil ⁵Instituto de Astrofísica de La Plata (UNLP-CONICET), La Plata, Argentina

⁶ Faculdad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Paseo del Bosque s/n, 1900, La Plata, Argentina

Submitted to ApJ

ABSTRACT

A spectral survey of methyl acetylene (CH_3CCH) was conducted toward the hot molecular core/outflow G331.512-0.103. Our APEX observations allowed the detection of 41 uncontaminated rotational lines of CH_3CCH in the frequency range between 172–356 GHz. Through an analysis under the local thermodynamic equilibrium assumption, by means of rotational diagrams, we determined T_{exc} =50±1 K, $N(CH_3CCH)$ =(7.5±0.4)×10¹⁵ cm⁻², $X[CH_3CCH/H_2]\approx (0.8-2.8)\times 10^{-8}$ and $X[CH_3CCH/CH_3OH] \approx 0.42 \pm 0.05$ for an extended emitting region (~10"). The relative intensities of the K=2 and K=3 lines within a given K-ladder are strongly negatively correlated to the transitions? upper J quantum-number (r=-0.84). Pure rotational spectra of CH₃CCH were simulated at different temperatures, in order to interpret this observation. The results indicate that the emission is characterized by a non-negligible temperature gradient with upper and lower limits of ~ 45 and ~ 60 K, respectively. Moreover, the line widths and peak velocities show an overall strong correlation with their rest frequencies, suggesting that the warmer gas is also associated with stronger turbulence effects. The K=0 transitions present a slightly different kinematic signature than the remaining lines, indicating that they might be tracing a different gas component. We speculate that this component is characterized by lower temperatures, and therefore larger sizes. Moreover, we predict and discuss the temporal evolution of the CH₃CCH abundance using a two-stage zero-dimensional model of the source constructed with the three-phase Nautilus gas-grain code.

Keywords: Methyl acetylene — Molecular outflows — Hot Molecular Cores — Astrochemistry

1. INTRODUCTION

Star-forming regions play a key role in building the complex inventory of chemical species detected in astronomical environments, which in turn serve as powerful tools to study their surroundings. Through the observation of molecular emission lines, it is possible to constrain both the physical conditions and the chemical evolution of circumstellar sources, shedding light on the formation processes that connect different evolutionary

Corresponding author: Julia C. Santos santos@strw.leidenuniv.nl stages of star-formation. Methyl acetylene (CH₃CCH), also known as propyne, is a particularly good temperature probe, and has therefore been widely explored toward various star-forming objects (e.g., Fontani et al. 2002; Bisschop et al. 2007; Fayolle et al. 2015; Giannetti et al. 2017; Andron et al. 2018; Calcutt et al. 2019).

Massive stars $(L > 10^3 L_{\odot}; M > 8M_{\odot})$ greatly affect their surroundings through their feedback and the production of heavy elements. The formation process of those stars, however, is still less well-understood in comparison to the low-mass counterparts. Observational disadvantages, such as their complex cluster environments $(n_{\star} \gtrsim 100 \text{ pc}^{-3})$ and large distances involved $(d \ge 1 \text{ kpc})$, together with a considerably shorter evolu-

tionary timescale ($t_{KH} \leq 10^4$ yr for O-type stars), substantially impair the development of a solid massive starformation paradigm (Garay & Lizano 1999; Zinnecker & Yorke 2007; Tan et al. 2014; Krumholz 2015; Silva et al. 2017; Rosen et al. 2020).

Despite the lack of a thoroughly defined evolutionary sequence, some precursors of high-mass stars are well established. One example are Hot Molecular Cores (HMCs), which are one of the first manifestations of massive star formation (Cesaroni 2005). They are characterized by relatively high temperatures (>100 K) and high densities $(n_{H_2} \sim 10^5 - 10^8 \text{ cm}^{-3})$, linked with a compact (<0.1 pc), luminous (>10⁴ L_{\odot}) and massive (~10- $1000 \, M_{\odot}$) molecular core (Heaton et al. 1989; Gomez et al. 1995; Cesaroni et al. 1998; Cesaroni 2005; Bonfand et al. 2019). HMCs are associated with a rich molecular emission spectrum, which carries information on their chemical and physical properties, as well as their morphology and evolutionary stage (e.g., Caselli et al. 1993; Comito et al. 2005; Herbst & van Dishoeck 2009; Allen et al. 2018; Bonfand et al. 2019; Jørgensen et al. 2020; Gieser et al. 2021).

1.1. The source: G331.512-0.103

G331.512-0.103 (henceforth G331) is a massive and energetic hot molecular core/outflow system. It is located within the millimeter source G331.5-0.1 (see Merello et al. 2013b and references therein), at the tangent point of the Norma spiral arm (~ 7.5 kpc) (Bronfman et al. 1985, 2008), and which consists of a singularly extended and luminous complex at the center of a Giant Molecular Cloud (GMC). The parent GMC is located at the peak region of the southern molecular and massive-star-formation rings (i.e., the peaks of the azimuthally averaged radial distributions of molecular gas and of regions of massive star formation, Bronfman et al. 2000). Likewise, it is one of the most massive and active star forming clouds in the southern Galaxy (García et al. 2014). Observational evidence indicates that a Young Massive Stellar Object (YMSO) at the center of G331 drives a powerful outflow with $\sim 55 M_{\odot}$ of mass and a momentum of $\sim 2.4 \times 10^3 \ M_{\odot} \ {\rm km \ s^{-1}}$ (Bronfman et al. 2008), with lobes closely aligned with the line of sight. Observations with the Atacama Large Millimeter/submillimeter Array (ALMA) of the SiO(8-7), $H^{13}CO^{+}(4-3)$, $HCO^{+}(4-3)$ and CO(3-2) transitions revealed an expanding bubble geometry driven by stellar winds, which probably arises from the protostar confined in a compact H II region of ~ 5 arcsec (projected size of ~ 0.2 pc) (Merello et al. 2013a). Its emission spectrum exhibits great chemical lavishness, from common radicals and carbon-chain molecules to prebiotic and complex organic species (Merello et al. 2013a,b; Mendoza et al. 2018; Duronea et al. 2019; Hervías-Caimapo et al. 2019; Canelo et al. 2021). All those factors make G331 an exquisite source for astrochemical investigations by means of spectral surveys in sub-mm wavelenghts.

1.2. The molecule: Methyl acetylene

More than 200 molecules have been hitherto detected toward the interstellar and circumstellar media (e.g., McGuire 2018). Methyl acetylene (CH₃CCH) has been detected in a wide variety of sources. It has been proven to be a reliable tracer of physical conditions, such as temperature and density, and has been extensively studied toward star-forming regions (e.g., Buhl & Snyder 1973; Lovas et al. 1976; Hollis et al. 1981; Kuiper et al. 1984; Taniguchi et al. 2018; Guzmán et al. 2018; Bøgelund et al. 2019; Calcutt et al. 2019). It has also been observed toward extragalactic sources, such as M 82, NGC 253, and NGC 1068 (Mauersberger et al. 1991; Qiu et al. 2020), and toward a planetary nebula (Schmidt & Ziurys 2019).

The rotational transitions of CH₃CCH are characterized by two quantum numbers, namely the total angular momentum (J) and its projection on the principal symmetry axis (K) (Townes & Schawlow 1975; Müller et al. 2000). Because it is a symmetric rotor, CH_3CCH presents many transitions that are closely spaced in frequency: the so-called K-ladders. The capability to observe many lines in the same bandwidth reduces calibration uncertainties and yields more precise predictions. Moreover, due to its small electric dipole moment $(\mu = 0.75 \text{ D})$ (Dubrulle et al. 1978), line thermalization occurs at densities as low as $\sim 10^4$ cm⁻³ (e.g., Bergin et al. 1994; Fontani et al. 2002; Molinari et al. 2016). Transitions with $\Delta K \neq 0$ are forbidden, and therefore the relative population of different K-ladders are dictated by collisions. As a result, CH₃CCH acts as an excellent temperature probe. Indeed, this molecule's temperature sensitivity makes it a reliable tracer of physical conditions and passive heating (Giannetti et al. 2017; Molinari et al. 2016).

Aside from observational advantages, the intrinsic spectroscopic properties of CH_3CCH can directly affect the observed spectra and therefore can give powerful insights on the physics of the emitting gas. Rotational spectroscopy is a conspicuously fruitful technique to infer information on the molecular species and its environment (e.g., Domenicano & Hargittai 1992; Winnewisser et al. 2003; Grubbs & Cooke 2010), and is nowadays highly assisted by quantum-chemistry (e.g., Puzzarini et al. 2012b,a; Cernicharo et al. 2015; McGuire et al. 2016; Cazzoli et al. 2016; Cerqueira et al. 2020; Santos et al. 2020). The rotational Hamiltonian is described in terms of the zeroth-order rotational constants (A, B, and C), and the higher-order centrifugal-distortion constants (e.g., D_J , D_{JK} , D_K , H_J , H_{JK} , H_{KJ} and H_K , for symmetric tops). These constants can be measured from laboratory experiments guided by theoretical simulations, which in turn play a fundamental role in line assignments of radioastronomical observations (e.g., Belloche et al. 2014; Coutens et al. 2016; Melli et al. 2018; Belloche et al. 2019).

In this work, we conducted a spectral survey of CH_3CCH toward G331, resulting in the detection of 41 uncontaminated lines. The goal of this work is to analyze the excitation conditions of methyl acetylene in this source, which gives information on the chemical and physical conditions of the gas and consequently helps unveiling the early stages of massive star-formation. Quantum-chemical properties associated with this molecule are especially relevant to explore the small-scale structure of the source. In section 2, we delineate the observational procedure. In section 3, we present our results on the line identifications and radiative analysis. Those results are then discussed in section 4. The observed abundance is discussed in the context of a chemical model of our source in section 5. Finally, our conclusions and perspectives are presented in section 6.

2. OBSERVATIONS

The observations were obtained with the Atacama Pathfinder EXperiment (APEX) telescope (Güsten et al. 2006) using the single-point mode toward the coordinates of the source (Bronfman et al. 2008) RA, DEC $(J2000) = 16^{h} 12^{m} 10.1^{s}, -51^{\circ} 28' 38.1''$. The Swedish-ESO PI Instrument (SEPIA180; Belitsky et al. 2018) was used, together with APEX-1 and APEX-2 receivers of the Swedish Heterodyne Facility Instrument (SHeFI; Vassilev et al. 2008), to observe nine frequency setups within the intervals 170-205 GHz and 222-307 GHz, respectively. As backend for the APEX-1 and APEX-2 receivers, the eXtended bandwidth Fast Fourier Transform Spectrometer2 (XFFTS2) was used, which consists of two units with 2.5 GHz bandwidth divided into 32768 channels each. SEPIA180 is a dual-polarization sideband-separated (2SB) receiver, which is able to observe two 4 GHz band, separated by 12 GHz, simultaneously. Thus, it covers 100% of the SEPIA ALMA Band 5 receiver channel IF band (159–211 GHz). The spectral resolution, corresponding to velocity resolutions, was obtained between 0.06 and 0.13 km s⁻¹ for a noise level of \sim 30–50 mK. While all analyses were performed with the original data, for further clarity and uniformity in their

graphical depiction, the resolution of the spectra exhibited in the present work were degraded to a common value of 1 km s⁻¹. The original intensity, obtained in a scale of antenna temperature corrected for atmospheric attenuation (T_A), was converted to the main-beam temperature (T_{mb}) scale using the main-beam efficiencies $\eta_{mb} = 0.80$ for SEPIA180¹, and $\eta_{mb} = 0.75$ and 0.73 for APEX-1 and APEX-2, respectively² (e.g. Canelo et al. 2021). The Half Power Beam Width (HPBW) values vary between ~17-39".³ We adopted a calibration uncertainty of about 10% (Dumke & Mac-Auliffe 2010).

The data reduction was carried out using the CLASS package of the GILDAS software⁴ (Pety et al. 2005; Gildas Team 2013). First-degree polynomial baselines were removed from each individual scan, which were subsequently averaged into a final spectrum. Line identifications were performed using the Weeds extension of CLASS, in combination with spectroscopic databases such as NIST⁵ Recommended Rest Frequencies for Observed Interstellar Molecular Microwave Transitions (Lovas 2004), CDMS⁶ (Müller et al. 2001, 2005; Endres et al. 2016), JPL⁷ (Pickett et al. 1998), and Splatalogue⁸. The establishment of a molecular identification warrants the satisfaction of a set of standard requirements (Snyder et al. 2005). Accordingly, the following criteria were used in order to confirm a detection: the peak frequencies of the observed lines should be consistent with the systemic velocity of G331 $(V_{lsr} \sim -90 \text{ km s}^{-1})$, the intensities of the observed lines should surpass the threshold of 3 rms noise, and lines predicted through Local Thermodynamic Equilibrium (LTE) modeling should agree with the observations. The radiative studies, including the optical depth estimation, were based on analyses carried out with the CASSIS software⁹ (Vastel et al. 2015)—assuming LTE. CASSIS has been developed by IRAP-UPS/CNRS.

3. RESULTS

3.1. Line analysis of CH₃CCH

We detected a total of 41 lines of CH_3CCH spread across the spectral band, from ~ 170.84 GHz to ~ 307.60

- ¹ http://www.apex-telescope.org/telescope/efficiency/
- 2 https://www.apex-telescope.org/telescope/efficiency/index.php. old
- $^{3} \ {\rm https://www.apex-telescope.org/instruments/}$
- $^{4}\;\rm https://www.iram.fr/IRAMFR/GILDAS/$
- ⁵ https://physics.nist.gov/cgi-bin/micro/table5/start.pl
- ⁶ https://cdms.astro.uni-koeln.de/
- ⁷ https://spec.jpl.nasa.gov/
- 8 https://www.cv.nrao.edu/php/splat/
- ⁹ http://cassis.irap.omp.eu/

GHz. Considering that CH_3CCH exhibits a K-ladder spectral signature, in Figure 1 we have displayed all the observed K-ladder structures from the J=10-9 to J=18-17 rotational levels.

Spectroscopic parameters obtained through Gaussian fittings to the lines are summarized in Table 1. The fittings were performed to the spectra at their full resolution of $\sim 0.06-0.13$ km s⁻¹.

For each observed J_u level, all transitions with K=3-0were detected. In some cases (see Table 1), lines with K=4 were not detectable above the 3 rms noise threshold, and were not considered in our analyses. Nevertheless, all K-ladders include a minimum of four detected transitions. The CH₃CCH lines present a low velocity dispersion (± 0.68 km s⁻¹) around the systemic velocity of the source and a mean systemic velocity of $\bar{V}_{lsr} = -90.0 \pm 0.3$ km s⁻¹, obtained through averaging the velocities of all 41 lines. Line profiles are narrow, with average values of 4.9 ± 0.8 km s⁻¹, and observed line widths also show an overall small dispersion, with FWHM values ranging from 4.15 to 5.68 km s⁻¹. This indicates that the emission originates from a quiescent region, as will be discussed in § 4.1.

3.2. Rotational diagrams and physical properties

In order to estimate the excitation temperature (T_{exc}) and column density (N) of CH₃CCH, rotational diagrams were constructed. Under the assumption of LTE, one can derive the T_{exc} and N of an optically thin emission that uniformly fills the antenna beam from:

$$\ln\left(\frac{N_u}{g_u}\right) = \ln\left(\frac{N}{Q(T_{exc})}\right) - \frac{E_u}{k_B T_{exc}} \tag{1}$$

where Q is the species' partition function at T_{exc} and k_B is the Boltzmann constant. N_u , g_u and E_u are, respectively, the column density, the statistical weight and the energy of the upper level (Goldsmith & Langer 1999). A plot of $\ln(N_u/g_u)$ versus E_u of the observed lines will thus correspond to a straight line, whose slope and yintercept are defined by $1/T_{exc}$ and $\ln(N/Q(\mathbf{T_{exc}}))$, respectively. For a spatially unresolved emitter, a beamdilution correction factor, $(\Delta\Omega_a/\Delta\Omega_s)$, should be introduced on the right-hand side of Equation (1). For a preliminary analysis, we adopted an average source size of 5", in accordance with previous observations of G331 (e.g. Hervías-Caimapo et al. 2019; Canelo et al. 2021, and references therein).

The rotational diagram of CH₃CCH constructed under the assumption of optically thin lines is displayed in blue in the upper panel of Figure 2. We derived $N(\text{CH}_3\text{CCH}) = (2.5\pm0.1) \times 10^{16} \text{ cm}^{-2}$ and $T_{exc} = 50\pm1$ K from the fit $(\chi^2_{red}=1.49)$. The uncertainties were computed following the formalism of the CASSIS software¹⁰, which takes into account the calibration error of 10%.

To explore the validity of the optically thin assumption, we estimated the lines' optical depths (τ) using CASSIS. The opacity correction is incorporated to Equation (1) through iterative calculations of the optical-depth correction factor C_{τ} (Goldsmith & Langer 1999), associated with the photon escape probability:

$$C_{\tau} = \frac{\tau}{1 - e^{-\tau}} \tag{2}$$

The optical-depth correction factor is introduced on the right-hand side of Equation (1). The iterations stop when a solution for T_{exc} and N converges. That is to say, when the difference between the last two iterations is smaller than 1%. The opacity-corrected rotational diagram of CH₃CCH is displayed in red in both panels of Figure 2. From the fit ($\chi^2_{red} = 2.57$) we obtained $N(\text{CH}_3\text{CCH}) = (4.5 \pm 0.3) \times 10^{16} \text{ cm}^{-2}$ and $T_{exc} = 42 \pm 1.0 \text{ K}$. This correction leads to a new scaling of the diagram's ordinate axis, resulting in a decrease of 17% and an increase of 80% in the values of T_{exc} and N, respectively.

Given the rather low temperatures derived from the rotational diagrams, we infer that the CH₃CCH emission is originated from a cooler gas in the outer envelope of the source, in accordance with other observational works toward star-forming regions (Churchwell & Hollis 1983; Nagy et al. 2015; Andron et al. 2018; Hervías-Caimapo et al. 2019). Thus, we expect CH₃CCH to trace a more extended region. The rotational diagram constructed for an extended source (size of 10") is displayed in the lower panel of Figure 2. From the fit under the optically thin assumption (χ^2_{red} =1.43), we derived $N(\text{CH}_3\text{CCH})=(6.9\pm0.4)\times10^{15} \text{ cm}^{-2}$ and $T_{exc}=52\pm1$ K.

After including the opacity correction factor, we derived $N(\text{CH}_3\text{CCH}) = (7.5 \pm 0.4) \times 10^{15} \text{ cm}^{-2}$ and $T_{exc} = 50 \pm 1 \text{ K}$ from the fit $(\chi^2_{red} = 1.51)$. These values were employed to model the emission of CH₃CCH under LTE conditions with WEEDS (see Figure 3). Overall, the synthetic spectra are in good agreement with the observations, although the accordance is sensibly higher for K-ladders with lower J quantum-numbers (see the discussion in § 4.2). For the 10" source size, the contribution of C_{τ} to the diagram is much less significant than for more compact emissions. Indeed, the corrected fit yielded T_{exc} and N values only slightly different from the optically thin scenario, with respective changes of -2.09% and 8.69%. Table 2 summarizes the excita-

¹⁰ http://cassis.irap.omp.eu/docs/RadiativeTransfer.pdf



Figure 1. Spectral lines of CH₃CCH identified toward G331 across the frequency range of 170.850–307.590 GHz. Each panel displays the K-ladder lines for the rotational transitions a) $J = 10_K - 9_K$, b) $J = 11_K - 10_K$, c) $J = 12_K - 11_K$, d) $J = 13_K - 12_K$, e) $J = 14_K - 13_K$, f) $J = 15_K - 14_K$, g) $J = 16_K - 15_K$, h) $J = 17_K - 16_K$ and i) $J = 18_K - 17_K$. The asterisk (*) indicates lines below the 3σ threshold. Linear baselines were subtracted from each spectrum, and the spectral resolution was resampled to a common channel of approximately 1 km s⁻¹.

tion temperatures and column densities derived from the rotational-diagram analyses.

In terms of the reduced χ^2 values, the fit of the 5" diagram with opacity corrections was less accurate in comparison with the uncorrected counterpart, whereas no significant change was observed for the 10" diagrams. This is a consequence of the underestimation of the source size in the 5" scenario, which is compensated by overestimating the lines' optical depth—in particular, for the K=0 and K=1 transitions at lower J values (see Figure 3). Indeed, the CH₃CCH emission seems to be optically thin (Churchwell & Hollis 1983; Fontani et al. 2002), especially for the range of J quantum-numbers observed in this work.

3.3. Relative intensities

The relative intensities of the lines within a K-ladder will change depending on the temperature of the environment. To explore this relation, the rotational spectra of CH₃CCH were simulated at temperatures ranging from 10 K to 100 K, using the PGOPHER general purpose software (Western 2016). The rotational constants (A and B), the quartic and sextic centrifugal distortion constants (D_J, D_{JK}, H_J, H_{JK} and H_{KJ}), and the dipole

Table 1. Spectral lines of CH₃CCH observed toward G331 and their parameters obtained from Gaussian fits. *Lines below the 3σ threshold.

Quantum Numbers		Frequency	E_u	A_u	$\int T_A dv$	V_{lsr}	FWHM		
J_u	K_u	$-J_l$	K_l	(MHz)	(K)	(10^{-5} s^{-1})	$(K \text{ km s}^{-1})$	$(\mathrm{km} \mathrm{s}^{-1})$	$(\mathrm{km}~\mathrm{s}^{-1})$
10	4	9	4	170853.576	160.7	1.43	0.24 ± 0.01	-90.02 ± 0.09	4.4 ± 0.2
10	3	9	3	170876.410	110.1	1.55	1.43 ± 0.01	-89.89 ± 0.02	4.57 ± 0.06
10	2	9	2	170892.726	74.0	1.63	1.75 ± 0.02	-89.77 ± 0.03	4.41 ± 0.06
10	1	9	1	170902.518	52.3	1.68	3.04 ± 0.06	-89.74 ± 0.04	4.4 ± 0.1
10	0	9	0	170905.783	45.1	1.70	3.95 ± 0.04	-89.34 ± 0.02	4.99 ± 0.06
11	4	10	4	187936.230	169.7	1.97	0.38 ± 0.01	-90.40 ± 0.08	4.76 ± 0.2
11	3	10	3	187961.342	119.2	2.10	1.87 ± 0.02	-89.96 ± 0.02	4.68 ± 0.05
11	2	10	2	187979.286	83.0	2.20	2.07 ± 0.02	-89.84 ± 0.02	4.45 ± 0.06
11	1	10	1	187990.055	61.4	2.25	3.53 ± 0.07	-89.83 ± 0.04	4.6 ± 0.1
11	0	10	0	187993.645	54.1	2.27	4.43 ± 0.04	-89.39 ± 0.02	5.06 ± 0.06
12	4	11	4	205018.114	179.6	2.63	0.48 ± 0.01	-89.89 ± 0.07	5.0 ± 0.2
12	3	11	3	205045.501	129.0	2.77	1.97 ± 0.02	-89.99 ± 0.02	4.80 ± 0.06
12	2	11	2	205065.070	92.9	2.88	2.04 ± 0.02	-89.86 ± 0.02	4.52 ± 0.05
12	1	11	1	205076.816	71.2	2.94	3.51 ± 0.06	-89.94 ± 0.04	4.6 ± 0.1
12	0	11	0	205080.732	64.0	2.96	4.46 ± 0.05	-89.43 ± 0.03	5.18 ± 0.07
13	4	12	4	222099.153^*	190.2	3.42	0.72 ± 0.04	-89.9 ± 0.2	6.6 ± 0.4
13	3	12	3	222128.815	139.7	3.57	1.97 ± 0.04	-90.04 ± 0.04	4.5 ± 0.1
13	2	12	2	222150.010	103.5	3.69	2.33 ± 0.03	-90.07 ± 0.03	4.84 ± 0.08
13	1	12	1	222162.730	81.9	3.75	3.74 ± 0.09	-90.15 ± 0.06	5.0 ± 0.1
13	0	12	0	222166.971	74.6	3.78	4.43 ± 0.06	-89.52 ± 0.03	5.24 ± 0.09
14	4	13	4	239179.281	201.7	4.34	0.44 ± 0.02	-90.62 ± 0.07	4.1 ± 0.2
14	3	13	3	239211.215	151.1	4.51	2.02 ± 0.03	-90.15 ± 0.03	4.55 ± 0.07
14	2	13	2	239234.034	115.0	4.63	2.08 ± 0.03	-90.09 ± 0.03	4.51 ± 0.06
14	1	13	1	239247.728	93.3	4.70	3.69 ± 0.07	-90.29 ± 0.05	5.2 ± 0.1
14	0	13	0	239252.294	86.1	4.73	4.25 ± 0.06	-89.50 ± 0.03	5.34 ± 0.09
15	4	14	4	256258.426^*	214.0	5.41	1.2 ± 0.1	-89.1 ± 0.3	7.3 ± 0.7
15	3	14	3	256292.630	163.4	5.59	2.29 ± 0.06	-90.14 ± 0.06	5.1 ± 0.1
15	2	14	2	256317.071	127.3	5.72	2.13 ± 0.06	-90.02 ± 0.06	4.9 ± 0.1
15	1	14	1	256331.739	105.7	5.80	3.23 ± 0.06	-90.14 ± 0.04	4.8 ± 0.1
15	0	14	0	256336.629	98.4	5.83	4.00 ± 0.05	-89.54 ± 0.04	5.43 ± 0.09
16	4	15	4	273336.519	227.1	6.64	0.58 ± 0.02	-90.30 ± 0.08	5.4 ± 0.2
16	3	15	3	273372.990	176.6	6.84	2.08 ± 0.02	-90.17 ± 0.03	4.91 ± 0.07
16	2	15	2	273399.051	140.4	6.97	2.13 ± 0.02	-90.16 ± 0.03	4.93 ± 0.07
16	1	15	1	273414.692	118.8	7.06	3.48 ± 0.06	-90.42 ± 0.04	5.5 ± 0.1
16	0	15	0	273419.906	111.5	7.09	3.82 ± 0.02	-89.66 ± 0.02	5.48 ± 0.09
17	4	16	4	290413.488^*	241.0	8.04	0.58 ± 0.02	-90.2 ± 0.1	5.3 ± 0.2
17	3	16	3	290452.224	190.5	8.25	2.40 ± 0.05	-90.47 ± 0.04	5.4 ± 0.1
17	2	16	2	290479.904	154.4	8.40	2.07 ± 0.03	-90.27 ± 0.03	4.9 ± 0.1
17	1	16	1	290496.516	132.7	8.49	3.53 ± 0.06	-90.68 ± 0.04	5.7 ± 0.1
17	0	16	0	290502.054	125.5	8.52	3.81 ± 0.05	-89.88 ± 0.04	5.4 ± 0.1
18	4	17	4	307489.264^*	255.8	9.62	0.52 ± 0.01	-90.13 ± 0.07	5.2 ± 0.2
18	3	17	3	307530.263	205.3	9.84	1.88 ± 0.02	-90.23 ± 0.03	4.90 ± 0.06
18	2	17	2	307559.559	169.1	10.00	1.73 ± 0.02	-90.23 ± 0.03	4.76 ± 0.07
18	1	17	1	307577.141	147.5	10.10	2.91 ± 0.04	-90.50 ± 0.03	5.39 ± 0.08
18	0	17	0	307583.003	140.3	10.10	3.27 ± 0.05	-89.76 ± 0.04	5.6 ± 0.1

moment (μ) employed in the simulations are listed in Table 3.

The transition intensities are given by (Western 2016):

Table 2. Values of T_{exc} and N of CH₃CCH toward G331 obtained from the analysis of the rotational diagrams.

Parameter	Source	e size: $5''$	Source size: 10"		
	With C_{τ}	Without \mathbf{C}_{τ}	With C_{τ}	Without C_{τ}	
T_{exc} [K]	42 ± 1	51 ± 1	50 ± 1	52 ± 1	
$N \ [imes 10^{16} \ { m cm}^{-2}]$	4.5 ± 0.3	2.5 ± 0.1	0.75 ± 0.04	0.69 ± 0.04	
χ^2_{red}	2.57	1.49	1.51	1.42	



Figure 2. Upper panel: Rotational diagram of the CH_3CCH lines for a source size of 5". Red circles and blue squares correspond to the lines with and without opacity corrections, respectively. Least squares fits to the points are also shown in their respective color. The error bars are propagated from the Gaussian fit to the lines and take into consideration a calibration uncertainty of 10% (Santos 2021) Lower panel: Same as the upper panel, but for a source size of 10" in diameter.

$$I = \frac{S}{Q(T)} \left[\exp\left(\frac{-E_l}{k_B T}\right) - \exp\left(\frac{-E_u}{\mathbf{k_B} T}\right) \right], \quad (3)$$

which is simply the line strength S times the Boltzmann factor, normalized to the partition function Q(T) at the given temperature. E_l and E_u correspond to the energies of the lower and the upper states, respectively. In Figure 4, we plot the intensity ratios I[K=n]/I[K=0] of the predicted lines within a K-ladder versus the temper-

Table 3. Spectroscopic parameters employed in the spectrum simulations. References: ^{*a*}JPL database; ^{*b*}(Dubrulle et al. 1978); ^{*c*}(Muenter & Laurie 1966).

Parameter	Value	Parameter	Value
A [MHz]	158590^{a}	H_J [Hz]	0.0097^{b}
B [MHz]	8545.87712^{b}	\mathbf{H}_{JK} [Hz]	0.935^{b}
D_J [kHz]	2.9423^{b}	\mathbf{H}_{KJ} [Hz]	5.23^{b}
D_{JK} [kHz]	163.423^{b}	μ [D]	0.75^{c}

ature, with n = 1, 2 and 3. We show the results for the J=10-9 and J=18-17 transitions, corresponding to the extremes of the observed bandwidth.

As seen in the plot, the behavior of the intensity ratios are overall analogous for the range of J values explored in this work. At low temperatures, the spectrum is dominated by the K=0 transitions, with small contributions from the other K levels. As the temperature rises, however, the ratios of all K transitions increase at different rates. The intensity ratios of both K=1and K=2 transitions, compared to K=0, follow a similar trend. They show an initial stage of rapid growth, which becomes gradually slower as the temperature continues to rise. As can be seen for the K=1 transitions, the curve will eventually reach a plateau, which stabilizes the K=1/K=0 intensity ratios at values around 0.8–0.9. Comparatively, the ratios for K=3 present a more dramatic increase. At around 50 K, the K=3 lines become more intense than the K=2 transitions. Further, at around 100 K, their intensities surpass those of the K=1 transitions. At even higher temperatures, they ultimately become the most intense line of the given Kladder.

This behavior can be explained by nuclear spin statistics: symmetric tops with three equivalent hydrogens, such as CH₃CCH, have two different spin symmetries namely, A and E. The A states correlate with the transitions with K=0, 3, 6, 9, ..., whereas the E states correspond to the remaining transitions, such as K=1, 2, 4, 5, ... (e.g. Strom et al. 2020). Those states have relative statistical weights of A:E = 2:1 (Herzberg 1945), and therefore the population distribution of the transitions at higher temperatures will favor the K=3 states. At lower temperatures, however, the energy acts as a limiting factor. This phenomenon makes the CH₃CCH K-



Figure 3. Spectra of CH₃CCH observed toward G331. Upper pannels: J=10-9, ~ 170.87–170.91 GHz; Lower pannels: J=18-17, ~ 307.52–307.59 GHz. Left pannels: LTE models simulated for a compact emission (5"). Right pannels: LTE models simulated for an extended emission (10"). For all panels, the black and red spectra correspond to the observations and the LTE models, respectively. The dashed line indicates the systemic velocity of the source. The abscissas of all spectra were adjusted to the systemic velocity of the K=0 transition.

ladder profile highly sensitive to the local temperature, in particular with regards to the relative intensities of the K=3 transitions.

As can be seen in Figure 1, we serendipitously observed a frequency window in which the CH₃CCH Kladder profiles gradually change as a function of the rotational quantum number. For low J values, which are associated with cooler regions, the K=2 lines are more intense than the K=3 (see panel a). However, as we observe transitions with higher J values, which are consequently associated with warmer regions, the K=2/K=3intensity ratios consistently decrease. From J=15-14onward, the K=3 lines become more intense than the K=2 (see panel i). In Figure 5, we plot the K=2/K=3ratios of the observed areas listed in Table 1 as a function of the upper J value. This plot yields a Pearson correlation coefficient of r=-0.84, which, given the small FWHM dispersion of the observed lines, clearly indicates a decreasing trend of the K=2/K=3 intensity ratios with J. This result strongly suggests that the lines

are tracing a region with a temperature gradient, as will be discussed in \S 4.2.

3.4. Search for the CH_3CCH isotopologues

The ¹³C isotopologues of CH₃CCH, as well as its deuterated forms, have been successfully detected toward multiple interstellar sources (Gerin et al. 1992; Markwick et al. 2002, 2005; Belloche et al. 2013; Potapov et al. 2016; Halfen et al. 2017; Schmidt & Ziurys 2019; Agúndez et al. 2019). Recently, Agundez et al. (2021) have detected the two doubly deuterated forms of methyl acetylene—CHD₂CCH and CH₂DCCD—toward the dense core L483. They derived abundance ratios of CH₃CCH/CHD₂CCH=34±10 and CH₃CCH/CH₂DCCD=42±13, which are only a few times less than the singly deuterated counterparts.

Motivated by the copious amount of CH₃CCH line detections in G331, we searched across the survey for the ¹³C and D isotopologues of CH₃CCH. Only the 10_0-9_0 and 10_1-9_1 lines of ¹³CH₃CCH were detected above the 3σ threshold. However, they are blended,



Figure 4. Intensity ratios I[K=n]/I[K=0] versus temperature for transitions within a given K-ladder. Purple squares represent ratios with n = 1, green circles represent ratios with n = 2, and red triangles represent ratios with n = 3. The solid and dashed lines correspond to, respectively, the J=10-9 and J=18-17 K-ladders. The red and blue regions represent upper and lower limits to the temperature profile of the source and are discussed in § 4.2.



Figure 5. Ratios of the areas for K=2 and K=3 lines versus the upper J value of the transition. Areas and uncertainties were obtained through Gaussian fittings of the observed data using CLASS. The dashed line indicates a ratio of unity.

and just marginally above the detection limit, and so we cannot reliably confirm these detections. Assuming a ratio of ${}^{12}C/{}^{13}C \sim 20$, as estimated for G331 and other sources toward the Galactic center (Wilson & Rood 1994; Requena-Torres et al. 2006; Mendoza et al. 2018; Duronea et al. 2019; Yan et al. 2019), we derive a ${}^{13}CH_3CCH$ column density of $N \sim 3.75 \times 10^{14}$ cm⁻². This column density was used to model the emission of the $10_K - 9_K$ K-ladder of ${}^{13}CH_3CCH$ under LTE condi-

tions (Figure 6), which is shown to be consistent with the observed spectrum of G331.



Figure 6. Modelled 10_K – 9_K K-ladder of ¹³CH₃CCH (in red) superimposed on the spectrum observed toward G331 (in black).

4. DISCUSSION

4.1. Origin of the emission

Hervías-Caimapo et al. (2019) have observed several molecular lines toward G331 and classified them according to the line profile, with narrow lines tracing the emission from the core ambient medium and lines with broad



Figure 7. Schematic view of the physical model proposed for G331. Based on the models of Merello et al. (2013b); Duronea et al. (2019); Hervías-Caimapo et al. (2019).

wings tracing the outflow and shocked region. Accordingly, we expect from the narrow line profiles that the observed CH_3CCH emission is originated from a quiescent (not expanding) core medium. This region is also associated with lower temperatures (below 100 K), which is consistent with the excitation temperatures derived from the rotational diagram. A schematic view of the physical model of G331, which considers the emission of CH_3CCH and other molecular tracers, is presented in Figure 7.

4.2. Physical conditions: gas temperatures

In § 3.4, we discussed the dependence of the K=2/K=3 intensity ratios of CH₃CCH with increasing temperatures. This inversion in relative populations within the same K-ladder is observed toward G331, and can be easily inferred by visual inspection (see Figure 1).

States with higher J quantum numbers require higher energies to be populated, and therefore tend to be associated with warmer regions than the lower J states. Given that the K=2/K=3 intensity ratios are negatively correlated with J (see Figure 5), we suggest that the CH₃CCH emission is in fact described by a temperature gradient, which explains the behavior seen in Figure 1. In order to estimate the extent of the temperature gradient, we have grouped the points in the rotational diagram into two subsets (Figure 8) separated at E_{up} ≈ 125 K—according to the behavior of the K=2/K=3intensity ratio of the K-ladder: the lower-energy group contains all K=2 transitions with $J \leq 14$, for which $K=2/K=3 \gtrsim 1$. Contrarily, the higher-energy subset encompasses all K=2 transitions with J > 14, for which $K=2/K=3 \leq 1$. The resulting T_{exc} and N obtained from the linear fit of each subset are listed in Table 4.



Figure 8. Rotational diagram of the CH_3CCH lines for a source size of 10". The points were divided into two subsets indicated by the dashed line. The points pertaining to the low- and high-energy subsets are shown in blue and red, respectively. Least squares fits to the points within each subset are shown. Circles and squares correspond to the lines with and without opacity corrections, respectively.

Assuming an optically thin scenario, the rotational diagram divided into two subsets yields temperature components of ~ 45 and ~ 60 K. Although the points in the rotational diagram show an overall fairly linear trend, the T_{exc} obtained from this analysis is highly sensitive to the fitted slope. Consequently, a difference of around 15 K arises between the derived values from the linear fittings of the two subsets. This temperature interval coincides with the region of Figure 4 where the intensity inversion of K=2 and K=3 occurs. Thus, one can regard the two temperature components extracted from the blue and red subsets in Figure 8 as, respectively, lower and upper limits to the temperature profile of the observed emitting region—indicated as the blue and red regions in Figure 4. The rotational diagram analysis of \S 3.2 is therefore suited to evaluate the large-scale emission of CH₃CCH, yielding an averaged temperature of $T_{exc} \sim 50.2$ K. The intrinsic spectroscopic properties of methyl acetylene, however, enabled us to perform a more direct and meticulous assessment of the temperature of the source. The same discussion is valid for the optically thick scenario.

The synthetic spectra under the LTE assumption shown in Figure 3 reproduces more accurately the observed data for lower J quantum-numbers. This gradual loss in accuracy for higher J transitions is likely a con-

Table 4. Values of T_{exc} and N of CH₃CCH obtained from the 2-component rotational diagrams for a source size of 10".

Parameter	Lower-energy	gy comp. (blue)	np. (blue) Higher-energy comp. (r		
	With C_{τ}	With C_{τ} Without C_{τ}		Without C_{τ}	
T_{exc} [K]	$45~\pm~2$	46 ± 2	60 ± 3	62 ± 3	
$N ~[imes 10^{15} ~{ m cm}^{-2}]$	7.9 ± 0.6	7.2 ± 0.5	5.7 ± 0.9	5.3 ± 0.8	
χ^2_{red}	0.87	0.82	1.19	1.14	

sequence of the temperature profile that leads to the intensity inversion around $J_u=14$ (see Figure 5). Indeed, throughout the entire ~170840–307600 MHz frequency range, modeled transitions with K=3 are less intense than the K=2 counterparts. Since using only one temperature component is not completely adequate to describe our observations, the model will be increasingly amiss as the bulk of the emission transits to regions with different average gas temperatures. Hence, our LTE model seems to best describe the cooler environment, traced by the lines with lower J values.

Previously, Hervías-Caimapo et al. (2019) have used ALMA to observed four CH₃CCH lines within the J=21-20 K-ladder toward G331, which showed a peak of emission located at a radius of ~1.2". From the rotational diagram constructed with the CH₃CCH lines, they obtained an excitation temperature of $T_{exc} = 70\pm7$ K. Their results are in accordance with our hypothesis of a temperature gradient, which indeed predicts that transitions with higher J quantum-numbers will be associated with higher excitation temperatures and more compact regions.

4.3. Kinematics

The spectroscopic parameters (see Table 1), obtained through fitting a Gaussian profile to each line, are directly related to the kinematics of the emitting region or, at least, of the gas where the bulk of the emission is originated. In Figure 9, we present a plot of the widths and peak velocities of the observed lines, which are listed in Table 1.

The line widths and peak velocities of transitions with the same quantum number K are overall strongly correlated. Indeed, the Pearson correlation coefficient of the K=0, K=1, K=2 and K=3 groups are, respectively, r=-0.85, r=-0.97, r=-0.78, and r=-0.80. This coefficient measures the linear correlation between two variants by means of the ratio of their covariance and the product of their standard deviation. Thus, $r=\pm 1$ correspond to perfect linear correlations (either positive or negative) and r=0 corresponds to no correlation. The lines with K=4 are fainter and harder to observe, and therefore were not included in the comparison. Their smaller signal-to-noise ratios hampers their analysis, which is reflected in their higher associated uncertainties reported in Table 1. Nonetheless, for each K group, there is a clear tendency for broader lines to present increasingly blueshifted velocities. The same trend was also observed by Di Francesco et al. (2004) for bright $(T_B^{\max} \ge 10 \sigma)$ N₂H⁺ lines toward the Ophiuchus A star-forming core: they found that the higher velocity lines were systematically broader, which they associate to infalling motions that increase local rotation speeds.

Interestingly, the K=0 lines are systematically broader and less blueshifted than the other groups, indicating that they might be tracing a slightly different region than the bulk of the emission of the other K transitions. At low temperatures, the contributions from transitions with $K \neq 0$ to the rotational spectrum of CH_3CCH are minor (see Figure 4), and the emission is dominated by the K=0 lines. As the temperature rises, the relative intensities of the $K \neq 0$ transitions rapidly increase, resulting in richer K-ladders. Thus, it is likely that the bulk of the K=0 emission is originated from a cooler and more extended region, which does not contribute as significantly to the lines with $K \neq 0$. These, in turn, must be tracing a hotter and more compact environment, resulting in a different kinematic signature. The systematic redshift of the peak velocities for the K=0 lines compared to the other groups also suggests interesting velocity features of the cold component. It is worth mentioning that the mean velocity shift between lines with the same quantum number K within the J=10-9 and J=18-17 K-ladders is ~ 0.49 km s⁻¹. which is more than three times the poorest obtained velocity resolution. Further surveys of the CH₃CCH emission toward G331 with higher angular resolution are imperative to better understand the different gas components, and the small-scale structure of the hot core.

In Figure 10, we show a comparison of the line widths versus the rest frequency, together with their Pearson coefficients. From the plots, it is clear that the FWHM is correlated to the line frequency, particularly for the lines with K=0 and K=1. Since thermal broadening effects are negligible for radio transitions at the considered temperature range (see, for example, the discussion of Fontani et al. 2002), the broader widths associated with higher J values indicate that the gas where the bulk of the emission is originated must become gradually more turbulent as the temperature increases. This is fairly



Figure 9. Line widths vs. peak velocities obtained from Gaussian fittings of the observed transitions. Blue crosses represent transitions with K=0, purple squares represent transitions with K=1, green circles represent transitions with K=2 and red triangles represent transitions with K=3. The black dashed line separates the blue- and redshifted regions relative to the adopted systemic velocity of G331 (-90 km s⁻¹).

reasonable, considering that the warmer emission comes from the inner parts of the envelope surrounding the Massive Young Stellar Object (MYSO).



Figure 10. Line widths vs. rest frequencies obtained from Gaussian fittings of the observed transitions with K=0, K=1, K=2 and K=3.

4.4. $CH_3 CCH$ abundances

The abundance of CH₃CCH relative to H₂ $(X(CH_3CCH))$ is defined as the ratio of the column densities $N(CH_3CCH)/N(H_2)$. We have estimated the H₂ column density in G331 from ALMA measurements of $N(H^{13}CO^+)\approx(1.5-3.2)\times10^{13}$ cm⁻² (Merello et al. 2013a,b; Hervías-Caimapo et al. 2019), adopting an abundance ratio $N(H^{13}CO^+)/N(H_2)=3.3\times10^{-11}$, as measured by Blake et al. (1987) in Orion KL. Additionally, a hydrogen column density of $N(H_2)=2.7\times10^{23}$ cm⁻² was reported for G331 from the continuum emission's surface density at 1.2 mm (Duronea et al. 2019). Considering those two limits for $N(H_2)$, we derived a ratio of $X(CH_3CCH)\approx(0.8-2.8)\times10^{-8}$.

In Table 5, we present a comparison of the CH_3CCH abundances with respect to H_2 in different massive starforming regions, including G331. Our derived abundance and excitation temperature agree well with the ones obtained for similar sources, which indicates similarities in the formation pathway of this molecule at such environments. The fact that this abundance comes mostly from the enveloping gas reinforces the conclusions drawn by other works (e.g., Öberg et al. 2013; Fayolle et al. 2015; Giannetti et al. 2017; Taniguchi et al. 2018) that the formation of methyl acetylene is not limited by heat input, and should take place during the earlier stages of star-formation. In other words, CH_3CCH is likely a so-called zeroth-generation molecule (Herbst & van Dishoeck 2009). Nonetheless, in the case of solidphase formation, heat will eventually be required in order for the molecule to desorb to the gas-phase.

We also estimate the fractional abundance of CH₃CCH with respect to methanol, $N(CH_3CCH)/N(CH_3OH)$, to be 0.42 ± 0.05. This value is listed in Table 5, together with the ratios obtained from the literature for other massive star-forming regions. The methanol column density in G331 is assumed to be of $\approx (1.8 \pm 0.2) \times 10^{16}$ cm⁻², as derived by Mendoza et al. (2018) under the LTE formalism and assuming the total CH_3OH abundance as the sum of the A-CH₃OH and E-CH₃OH contributions. This methanol emission, however, is thought to originate from a region of only 5''.3, which is considerably more compact than the emitting region of CH_3CCH . Given that the envelope abundance of methanol in G331 is not fully known, the derived $N(CH_3CCH)/N(CH_3OH)$ ratio is likely artificially lower than in reality. Nonetheless, it is still fairly close to the ratios derived in other sources, further strengthening the conclusion that they present similar chemistry.

5. CHEMICAL MODELING

Formation pathways of CH₃CCH in the ISM include gas-phase ion-neutral routes with $C_2H_2^+$ as a precursor (Schiff & Bohme 1979; Millar & Freeman 1984), neutralneutral reactions such as $CH + C_2H_4 \rightarrow CH_3CCH +$ H (Turner et al. 1999), as well as dissociative recombination reactions involving larger hydrocarbons (Calcutt et al. 2019). Grain-surface reactions are also proposed in order to explain the observed abundances of CH₃CCH (Hickson et al. 2016; Guzmán et al. 2018). Regarding massive environments, Taniguchi et al. (2019) have constructed hot-core models to investigate the formation pathways of cyanopolyynes and other carbonchain species, including CH₃CCH and c-C₃H₂, around MYSOs. They found chemical similarities between methyl acetylene, methane and cyanopolyynes, being all triggered by CH₄ sublimation from dust grains. Furthermore, those species are also shown to accumulate in the bulk of the ice until the temperature reaches their respective sublimation points. Comparatively, small reactive hydrocarbons such as CCH and CCS are shown not to accumulate onto dust grains, being readily destroyed in the gas-phase instead. Complementary, Andron et al. (2018) discuss the formation routes of methyl acetylene in the solar-type protostar IRAS 16293-2422. They show that, at large distances form the central star, the abundance of CH₃CCH in the solid phase is higher than in

the gas-phase, which indicates that it is efficiently produced on dust grains at low temperatures through successive hydrogenation of C_3 . Moreover, their models also indicates that reactions in the gas-phase with C_2H_4 and C_3H_5 as precursors can also effectively form methyl acetylene in the outer envelope.

In order to investigate the chemical evolution of CH_3CCH in G331, we constructed a time-dependent chemical model of the source using the three-phase gasgrain Nautilus code (Ruaud et al. 2016), in which the grain surface chemistry is distinguished from the bulk of the ice. The model consists of two zero-dimensional steps (i.e., the physical properties of the source are uniform and static), comprising the dark-cloud and the hotcore phases of star formation. The 1^{st} step, representing an initial dark-cloud phase, was computed using the elemental abundances listed in Table 6, as was computed in previous works about the chemistry of HNCO in G331 (Canelo et al. 2021). For the physical parameters, we adopted a gas temperature and density of 10 K and 1 \times 10^4 cm^{-3} , respectively. Then, we employed the abundances derived from the dark-cloud model after 10^5 yr as initial parameters for the 2^{nd} step: a "rapid" hot-core phase. For this second period we used T = 80 K and tested four different values for n_{H_2} : 1×10^6 , 5×10^6 , 1×10^7 and 5×10^7 cm⁻³. The gas and dust temperatures were set as equal throughout the entire simulation, as they have been shown to be coupled in dense sources (Merello et al. 2019). Moreover, standard visual extinction $(A_V = 10 \text{ mag})$ and cosmic ray ionization rates $(\zeta = 1.3 \times 10^{-17} \text{ cm}^{-3})$ were used. The standard chemical network presented in the KInetic Database for Astrochemistry (KIDA)¹¹ (Wakelam et al. 2015) catalog was employed in the simulations.

The temporal-evolution of the CH₃CCH abundance derived from the hot-core model for the different hydrogen densitites is presented in Figure 11. The abundance derived from our observations is best predicted by the model with $n_{H_2} = 5 \times 10^6$ at timescales as early as ~ 10³ yr, in agreement with the expected age of this source (Merello et al. 2013a). This is consistent with an extended and thus less dense emitting gas, as was inferred from the radiative analyses. Furthermore, the steady-state plateau in abundance reached after ~ 10³ yr is consistent with the hypothesis of CH₃CCH accumulating onto dust grains and enriching the gasphase upon desorption. Experiments on interstellar ice analogues exposed to energetic electrons ensued the formation of CH₃CCH, among other related molecules (Ab-

¹¹ http://kida.obs.u-bordeaux1.fr/

Table 5. Abundances of CH₃CCH relative to H₂ and CH₃OH observed toward high-mass star-forming sites. References: a(Taniguchi et al. 2018), b(Giannetti et al. 2017), c(Fayolle et al. 2015). *Derived from the average abundances reported by Giannetti et al. (2017). n.p. = not provided. The excitation temperatures are given in units of K.

Source	$/N({ m H_2})$	$/N(CH_3OH)$	$\mathbf{T}_{exc}^{\mathrm{CH}_{3}\mathrm{CCH}}$
$G12.89 + 0.49^{a}$	$4.2^{+2.8}_{-2.2} \times 10^{-8}$	$0.34_{-0.15}^{+0.28}$	33^{+20}_{-9}
$G16.86-2.16^{a}$	$3.2^{+1.9}_{-1.5} \times 10^{-8}$	$0.36\substack{+0.28\\-0.18}$	29^{+15}_{-8}
$G28.28-0.36^{a}$	$7.6^{+5.2}_{-4.4} imes 10^{-8}$	$1.61^{+1.4}_{-0.86}$	23^{+9}_{-6}
$\operatorname{ATLASGAL}^{b}$	$(0.5-2.5) \times 10^{-8}$	0.31^{*}	$34.5^{+24.6}_{-10.5}$
NGC 7538 $\mathrm{IRS9}^c$	n.p.	1.3 ± 0.4	47 ± 5
W3 IRS5^{c}	n.p.	2.2 ± 0.7	58 ± 8
$AFGL490^{c}$	n.p.	1.8 ± 0.8	41 ± 7
G331 (this work)	$(0.8 - 2.8) \times 10^{-8}$	0.42 ± 0.05	50 ± 1

Table 6. Initial gas-phase elemental abundances, with the format a(b) representing $a \times 10^b$. References: 1. Wakelam & Herbst (2008); 2. Jenkins (2009); 3. Hincelin et al. (2011); 4. low-metal abundances from Graedel et al. (1982); 5. depleted value from Neufeld et al. (2005).

Element	n_i/n_H^a	Ref.	Element	n_i/n_H^a	Ref.
H_2	0.5		He	9.0(-2)	1
Ν	6.2(-5)	2	0	2.4(-4)	3
C^+	1.7(-4)	2	S^+	1.5(-5)	2
Fe^+	3.0(-9)	4	Si^+	8.0(-9)	4
Na^+	2.0(-9)	4	Mg^+	7.0(-9)	4
Cl^+	1.0(-9)	4	\mathbf{P}^+	2.0(-10)	4
F	6.7(-9)	5			

planalp et al. 2019), which supports this hypothesis. Complementary observations of different species with related chemical networks are desirable to build a comprehensive view on the properties of the source, and will be included in future works focused on constructing a thorough chemical and physical model of G331.

6. CONCLUSIONS

We have conducted a spectral survey of CH₃CCH toward the Hot Molecular Core G331.512-0.103, resulting in the detection of 41 lines without contamination. To the best of our knowledge, this was the first spectral survey of methyl acetylene toward the source, resulting in interesting insights on the physics of the source. The spectral analysis was performed through rotational diagrams, assuming LTE, from which we derived an averaged excitation temperature of ~ 50 K for an extended emission. Thus, the bulk of the CH₃CCH emission is likely originated from a warm and extended gas, associated with the ambient core region of G331. Moreover, we obtained $N(CH_3CCH) = 7.5 \times 10^{15} \text{ cm}^{-2}$, $X[CH_3CCH/H_2] \approx (0.77 - 2.8) \times 10^{-8}$ and $X[CH_3CCH/CH_3OH] \approx 0.42 \pm 0.05$ from the observations, which are consistent with other single-dish



Figure 11. Abundance of CH_3CCH as a function of time, obtained from the hot-core model for different initial densities of H_2 . The gray area indicates the methyl-acetylene abundance derived from the observations.

observational works toward massive star-forming regions and suggest that CH_3CCH is a zeroth-generation molecule with similar chemistry throughout these environments.

The K=2/K=3 line-intensity ratios of transitions within a given K-ladder are strongly negatively correlated with J_u , which firmly suggests that the emission is arising from a region with a temperature gradient. An analysis of the rotational diagram separated into two subsections provides upper and lower limits of, respectively, ~60 and ~45 K for the temperature gradient. This approach enables us to assess the small-scale structure of the source with data from single-dish facilities. For a thorough analysis of the temperature profile, however, it is imperative to observe a wide range of CH₃CCH lines with a high angular resolution, by means of interferometry instruments such as ALMA.

The line widths of transitions with the same K quantum number are strongly correlated to their rest frequencies and peak velocities. Thus, we infer that the warmer regions, traced by the higher-frequency lines, are also associated with higher turbulence effects. Moreover, the K=0 transitions seem to be tracing a slightly different region than the rest of the CH₃CCH lines, as indicated by its different kinematic signature. Surveys of this species with higher angular resolution are also fundamental to further explore the origin of this differentiation.

A chemical model of CH₃CCH in G331, comprising of a dark-cloud phase followed by a hot-core collapse, predicts CH₃CCH abundances that agree with our observations for timescales of $t \gtrsim 10^3$ yr and densities of $n_{H_2} \sim 5 \times 10^6$ cm⁻³. This is in line with the expected age and size of this source.

ACKNOWLEDGMENTS

This publication is based on data acquired with the Atacama Pathfinder Experiment (APEX) under programme IDs C-094.F-9709B-2014, C-097.F-9710A-2016, C-099.F-9702A- 2017 and C-0102.F-9702B-2018. APEX is a collaboration between the Max-Planck-Institut fur Radioastronomie, the European Southern Observatory, and the Onsala Space Observatory. We thank the APEX staff for their helping during the observations and the anonymous referee for their constructive criticism that definitely improved this work. L.B. and R.F. acknowledge support from ANID project Basal AFB-170002. EM acknowledges support from the Brazilian agencies FAPESP, grant 2014/22095-6, and CNPq, grant 150465/2019-0. M.M. acknowledges support from ANID, Programa de Astronomía - Fondo ALMA-CONICYT, project 3119AS0001. RF acknowledge the support of ANID through the ALMA-CONICYT project 31180005. Finally, we would like to acknowledge Dr. Colin Western, who sadly passed away recently, for creating the PGOPHER Software. His contributions to the scientific community will remain evermore.

Facilities: Atacama Pathfinder EXperiment, APEX telescope.

Software: CASSIS (Vastel et al. 2015), GILDAS (Pety et al. 2005; Gildas Team 2013), NAUTILUS (Ruaud et al. 2016), PGOPHER (Western 2016).

REFERENCES

- Abplanalp, M. J., Góbi, S., & Kaiser, R. I. 2019, Physical Chemistry Chemical Physics (Incorporating Faraday Transactions), 21, 5378, doi: 10.1039/C8CP03921F
- Agúndez, M., Marcelino, N., Cernicharo, J., Roueff, E., & Tafalla, M. 2019, A&A, 625, A147, doi: 10.1051/0004-6361/201935164
- Agundez, M., Roueff, E., Cabezas, C., Cernicharo, J., & Marcelino, N. 2021, arXiv e-prints, arXiv:2104.04374. https://arxiv.org/abs/2104.04374
- Allen, V., van der Tak, F. F. S., & Walsh, C. 2018, A&A, 616, A67, doi: 10.1051/0004-6361/201732553
- Andron, I., Gratier, P., Majumdar, L., et al. 2018, MNRAS, 481, 5651, doi: 10.1093/mnras/sty2680
- Belitsky, V., Lapkin, I., Fredrixon, M., et al. 2018, A&A, 612, A23, doi: 10.1051/0004-6361/201731458
- Belloche, A., Garrod, R. T., Müller, H. S. P., & Menten, K. M. 2014, Science, 345, 1584, doi: 10.1126/science.1256678

- Belloche, A., Garrod, R. T., Müller, H. S. P., et al. 2019, A&A, 628, A10, doi: 10.1051/0004-6361/201935428
- Belloche, A., Müller, H. S. P., Menten, K. M., Schilke, P., & Comito, C. 2013, A&A, 559, A47, doi: 10.1051/0004-6361/201321096
- Bergin, E. A., Goldsmith, P. F., Snell, R. L., & Ungerechts, H. 1994, ApJ, 431, 674, doi: 10.1086/174518
- Bisschop, S. E., Jørgensen, J. K., van Dishoeck, E. F., & de Wachter, E. B. M. 2007, A&A, 465, 913, doi: 10.1051/0004-6361:20065963
- Blake, G. A., Sutton, E. C., Masson, C. R., & Phillips, T. G. 1987, ApJ, 315, 621, doi: 10.1086/165165
- Bøgelund, E. G., Barr, A. G., Taquet, V., et al. 2019, A&A, 628, A2, doi: 10.1051/0004-6361/201834527
- Bonfand, M., Belloche, A., Garrod, R. T., et al. 2019, A&A, 628, A27, doi: 10.1051/0004-6361/201935523
- Bronfman, L., Casassus, S., May, J., & Nyman, L. Å. 2000, A&A, 358, 521. https://arxiv.org/abs/astro-ph/0006104

- Bronfman, L., Cohen, R. S., Thaddeus, P., & Alvarez, H.1985, in The Milky Way Galaxy, ed. H. van Woerden,R. J. Allen, & W. B. Burton, Vol. 106, 331
- Bronfman, L., Garay, G., Merello, M., et al. 2008, ApJ, 672, 391, doi: 10.1086/522487
- Buhl, D., & Snyder, L. E. 1973, in Molecules in the Galactic Environment, ed. M. A. Gordon & L. E. Snyder, 187
- Calcutt, H., Willis, E. R., Jørgensen, J. K., et al. 2019, A&A, 631, A137, doi: 10.1051/0004-6361/201936323
- Canelo, C. M., Bronfman, L., Mendoza, E., et al. 2021, MNRAS, 504, 4428, doi: 10.1093/mnras/stab1163
- Caselli, P., Hasegawa, T. I., & Herbst, E. 1993, ApJ, 408, 548, doi: 10.1086/172612
- Cazzoli, G., Lattanzi, V., Kirsch, T., et al. 2016, A&A, 591, A126, doi: 10.1051/0004-6361/201628745
- Cernicharo, J., McCarthy, M. C., Gottlieb, C. A., et al. 2015, ApJL, 806, L3, doi: 10.1088/2041-8205/806/1/L3
- Cerqueira, H. B. A., Santos, J. C., Fantuzzi, F., et al. 2020, The Journal of Physical Chemistry A, 124, 6845, doi: 10.1021/acs.jpca.0c03529
- Cesaroni, R. 2005, in Massive Star Birth: A Crossroads of Astrophysics, ed. R. Cesaroni, M. Felli, E. Churchwell, & M. Walmsley, Vol. 227, 59–69, doi: 10.1017/S1743921305004369
- Cesaroni, R., Hofner, P., Walmsley, C. M., & Churchwell, E. 1998, A&A, 331, 709
- Churchwell, E., & Hollis, J. M. 1983, ApJ, 272, 591, doi: 10.1086/161322
- Comito, C., Schilke, P., Phillips, T. G., et al. 2005, ApJS, 156, 127, doi: 10.1086/425996
- Coutens, A., Jørgensen, J. K., van der Wiel, M. H. D., et al. 2016, A&A, 590, L6, doi: 10.1051/0004-6361/201628612
- Di Francesco, J., André, P., & Myers, P. C. 2004, ApJ, 617, 425, doi: 10.1086/425264
- Domenicano, A., & Hargittai, I. 1992, Accurate Molecular Structures: Their Determination and Importance (Oxford University Press)
- Dubrulle, A., Boucher, D., Burie, J., & Demaison, J. 1978, Journal of Molecular Spectroscopy, 72, 158, doi: https://doi.org/10.1016/0022-2852(78)90051-6
- Dumke, M., & Mac-Auliffe, F. 2010, in Proc. SPIE, Vol. 7737, Observatory Operations: Strategies, Processes, and Systems III, 77371J, doi: 10.1117/12.858020
- Duronea, N. U., Bronfman, L., Mendoza, E., et al. 2019, MNRAS, 489, 1519, doi: 10.1093/mnras/stz2087
- Endres, C. P., Schlemmer, S., Schilke, P., Stutzki, J., & Müller, H. S. P. 2016, Journal of Molecular Spectroscopy, 327, 95, doi: 10.1016/j.jms.2016.03.005

- Fayolle, E. C., Öberg, K. I., Garrod, R. T., van Dishoeck,
 E. F., & Bisschop, S. E. 2015, A&A, 576, A45,
 doi: 10.1051/0004-6361/201323114
- Fontani, F., Cesaroni, R., Caselli, P., & Olmi, L. 2002, A&A, 389, 603, doi: 10.1051/0004-6361:20020579
- Garay, G., & Lizano, S. 1999, PASP, 111, 1049, doi: 10.1086/316416
- García, P., Bronfman, L., Nyman, L.-Å., Dame, T. M., & Luna, A. 2014, ApJS, 212, 2, doi: 10.1088/0067-0049/212/1/2
- Gerin, M., Combes, F., Wlodarczak, G., Encrenaz, P., & Laurent, C. 1992, A&A, 253, L29
- Giannetti, A., Leurini, S., Wyrowski, F., et al. 2017, A&A, 603, A33, doi: 10.1051/0004-6361/201630048
- Gieser, C., Beuther, H., Semenov, D., et al. 2021, A&A, 648, A66, doi: 10.1051/0004-6361/202039670
- Gildas Team. 2013, GILDAS: Grenoble Image and Line Data Analysis Software. http://ascl.net/1305.010
- Goldsmith, P. F., & Langer, W. D. 1999, ApJ, 517, 209, doi: 10.1086/307195
- Gomez, Y., Garay, G., & Lizano, S. 1995, ApJ, 453, 727, doi: 10.1086/176434
- Graedel, T. E., Langer, W. D., & Frerking, M. A. 1982, ApJS, 48, 321, doi: 10.1086/190780
- Grubbs, G., & Cooke, S. 2010, Journal of Molecular Spectroscopy, 259, 120 ,
 - doi: https://doi.org/10.1016/j.jms.2009.12.003
- Güsten, R., Nyman, L. Å., Schilke, P., et al. 2006, A&A, 454, L13, doi: 10.1051/0004-6361:20065420
- Guzmán, A. E., Guzmán, V. V., Garay, G., Bronfman, L., & Hechenleitner, F. 2018, ApJS, 236, 45, doi: 10.3847/1538-4365/aac01d
- Halfen, D. T., Woolf, N. J., & Ziurys, L. M. 2017, ApJ, 845, 158, doi: 10.3847/1538-4357/aa816b
- Heaton, B. D., Little, L. T., & Bishop, I. S. 1989, A&A, 213, 148
- Herbst, E., & van Dishoeck, E. F. 2009, ARA&A, 47, 427, doi: 10.1146/annurev-astro-082708-101654
- Hervías-Caimapo, C., Merello, M., Bronfman, L., et al. 2019, ApJ, 872, 200, doi: 10.3847/1538-4357/aaf9ac
- Herzberg, G. 1945, Molecular Spectra And Molecular Structure. Volume II (D. Van Nostrand Company, Inc.)
- Hickson, K. M., Wakelam, V., & Loison, J.-C. 2016, Molecular Astrophysics, 3, 1, doi: 10.1016/j.molap.2016.03.001
- Hincelin, U., Wakelam, V., Hersant, F., et al. 2011, A&A, 530, A61, doi: 10.1051/0004-6361/201016328
- Hollis, J. M., Snyder, L. E., Blake, D. H., et al. 1981, ApJ, 251, 541, doi: 10.1086/159495

- Jenkins, E. B. 2009, ApJ, 700, 1299, doi: 10.1088/0004-637X/700/2/1299
- Jørgensen, J. K., Belloche, A., & Garrod, R. T. 2020, ARA&A, 58, 727,
- doi: 10.1146/annurev-astro-032620-021927

Krumholz, M. R. 2015, The Formation of Very Massive Stars, ed. J. S. Vink, Vol. 412, 43, doi: 10.1007/978-3-319-09596-7_3

- Kuiper, T. B. H., Kuiper, E. N. R., Dickinson, D. F., Turner, B. E., & Zuckerman, B. 1984, ApJ, 276, 211, doi: 10.1086/161604
- Lovas, F. J. 2004, Journal of Physical and Chemical Reference Data, 33, 177, doi: 10.1063/1.1633275
- Lovas, F. J., Johnson, D. R., Buhl, D., & Snyder, L. E. 1976, ApJ, 209, 770, doi: 10.1086/154774
- Markwick, A. J., Charnley, S. B., Butner, H. M., & Millar, T. J. 2005, ApJL, 627, L117, doi: 10.1086/432415
- Markwick, A. J., Millar, T. J., & Charnley, S. B. 2002, A&A, 381, 560, doi: 10.1051/0004-6361:20011381
- Mauersberger, R., Henkel, C., Walmsley, C. M., Sage, L. J., & Wiklind, T. 1991, A&A, 247, 307
- McGuire, B. A. 2018, ApJS, 239, 17, doi: 10.3847/1538-4365/aae5d2
- McGuire, B. A., Martin-Drumel, M.-A., Thorwirth, S., et al. 2016, Phys. Chem. Chem. Phys., 18, 22693, doi: 10.1039/C6CP03871A
- Melli, A., Melosso, M., Tasinato, N., et al. 2018, ApJ, 855, 123, doi: 10.3847/1538-4357/aaa899
- Mendoza, E., Bronfman, L., Duronea, N. U., et al. 2018, ApJ, 853, 152, doi: 10.3847/1538-4357/aaalec
- Merello, M., Bronfman, L., Garay, G., et al. 2013a, ApJL, 774, L7, doi: 10.1088/2041-8205/774/1/L7
- --. 2013b, ApJ, 774, 38, doi: 10.1088/0004-637X/774/1/38
- Merello, M., Molinari, S., Rygl, K. L. J., et al. 2019, MNRAS, 483, 5355, doi: 10.1093/mnras/sty3453
- Millar, T. J., & Freeman, A. 1984, MNRAS, 207, 405, doi: 10.1093/mnras/207.3.405
- Molinari, S., Merello, M., Elia, D., et al. 2016, ApJL, 826, L8, doi: 10.3847/2041-8205/826/1/L8
- Muenter, J. S., & Laurie, V. W. 1966, The Journal of Chemical Physics, 45, 855, doi: 10.1063/1.1727694
- Müller, H. S. P., Schlöder, F., Stutzki, J., & Winnewisser, G. 2005, Journal of Molecular Structure, 742, 215, doi: 10.1016/j.molstruc.2005.01.027
- Müller, H. S. P., Thorwirth, S., Bizzocchi, L., & Winnewisser, G. 2000, Zeitschrift Naturforschung Teil A, 55, 491, doi: 10.1515/zna-2000-0503
- Müller, H. S. P., Thorwirth, S., Roth, D. A., & Winnewisser, G. 2001, A&A, 370, L49, doi: 10.1051/0004-6361:20010367

- Nagy, Z., van der Tak, F. F. S., Fuller, G. A., & Plume, R. 2015, A&A, 577, A127, doi: 10.1051/0004-6361/201425014 Neufeld, D. A., Wolfire, M. G., & Schilke, P. 2005, ApJ, 628, 260, doi: 10.1086/430663 Öberg, K. I., Boamah, M. D., Fayolle, E. C., et al. 2013, ApJ, 771, 95, doi: 10.1088/0004-637X/771/2/95 Pety, J., Teyssier, D., Fossé, D., et al. 2005, A&A, 435, 885, doi: 10.1051/0004-6361:20041170 Pickett, H. M., Poynter, R. L., Cohen, E. A., et al. 1998, JQSRT, 60, 883, doi: 10.1016/S0022-4073(98)00091-0 Potapov, A., Sánchez-Monge, Á., Schilke, P., et al. 2016, A&A, 594, A117, doi: 10.1051/0004-6361/201628426 Puzzarini, C., Cazzoli, G., & Gauss, J. 2012a, JChPh, 137, 154311, doi: 10.1063/1.4758316 Puzzarini, C., Cazzoli, G., López, J. C., et al. 2012b, JChPh, 137, 024310, doi: 10.1063/1.4731284
- Qiu, J., Zhang, J., Zhang, Y., Jia, L., & Tang, X. 2020, A&A, 634, A125, doi: 10.1051/0004-6361/201935800
- Requena-Torres, M. A., Martín-Pintado, J., Rodríguez-Franco, A., et al. 2006, A&A, 455, 971, doi: 10.1051/0004-6361:20065190
- Rosen, A. L., Offner, S. S. R., Sadavoy, S. I., et al. 2020, SSRv, 216, 62, doi: 10.1007/s11214-020-00688-5
- Ruaud, M., Wakelam, V., & Hersant, F. 2016, MNRAS, 459, 3756, doi: 10.1093/mnras/stw887
- Santos, J. C. 2021, Master's thesis, University of Sao Paulo
- Santos, J. C., Rocha, A. B., & Oliveira, R. R. 2020, Journal of Molecular Modeling, 26, doi: 10.1007/s00894-020-04523-0
- Schiff, H. I., & Bohme, D. K. 1979, ApJ, 232, 740, doi: 10.1086/157334
- Schmidt, D. R., & Ziurys, L. M. 2019, ApJL, 881, L38, doi: 10.3847/2041-8213/ab3663
- Silva, A., Zhang, Q., Sanhueza, P., et al. 2017, ApJ, 847, 87, doi: 10.3847/1538-4357/aa88c6
- Snyder, L. E., Lovas, F. J., Hollis, J. M., et al. 2005, ApJ, 619, 914, doi: 10.1086/426677
- Strom, A. I., Gutiérrez-Quintanilla, A., Chevalier, M., et al. 2020, The Journal of Physical Chemistry A, 124, 4471, doi: 10.1021/acs.jpca.0c02900
- Tan, J. C., Beltrán, M. T., Caselli, P., et al. 2014, in Protostars and Planets VI, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning, 149, doi: 10.2458/azu_uapress_9780816531240-ch007
- Taniguchi, K., Herbst, E., Caselli, P., et al. 2019, ApJ, 881, 57, doi: 10.3847/1538-4357/ab2d9e
- Taniguchi, K., Saito, M., Majumdar, L., et al. 2018, ApJ, 866, 150, doi: 10.3847/1538-4357/aade97

- Townes, C. H., & Schawlow, A. L. 1975, Microwave Spectroscopy (Dover Publications)
- Turner, B. E., Terzieva, R., & Herbst, E. 1999, ApJ, 518, 699, doi: 10.1086/307300
- Vassilev, V., Meledin, D., Lapkin, I., et al. 2008, A&A, 490, 1157, doi: 10.1051/0004-6361:200810459
- Vastel, C., Bottinelli, S., Caux, E., Glorian, J. M., & Boiziot, M. 2015, in SF2A-2015: Proceedings of the Annual meeting of the French Society of Astronomy and Astrophysics, 313–316
- Wakelam, V., & Herbst, E. 2008, ApJ, 680, 371, doi: 10.1086/587734

- Wakelam, V., Loison, J. C., Herbst, E., et al. 2015, ApJS, 217, 20, doi: 10.1088/0067-0049/217/2/20
- Western, C. M. 2016, J. Quant. Spectrosc. Radiat. Transf., 186, 221
- Wilson, T. L., & Rood, R. 1994, ARA&A, 32, 191, doi: 10.1146/annurev.aa.32.090194.001203
- Winnewisser, G., Lewen, F., Thorwirth, S., et al. 2003, Chemistry – A European Journal, 9, 5501, doi: 10.1002/chem.200305192
- Yan, Y. T., Zhang, J. S., Henkel, C., et al. 2019, ApJ, 877, 154, doi: 10.3847/1538-4357/ab17d6
- Zinnecker, H., & Yorke, H. W. 2007, ARA&A, 45, 481, doi: 10.1146/annurev.astro.44.051905.092549